

# Two Spanning Disjoint Paths with Required Length in Augmented Cubes

Chung-Meng Lee

Department of Computer Science  
National Chiao Tung University  
cmlee@pu.edu.tw

Yuan-Hsiang Teng

Department of Computer Science  
National Chiao Tung University  
gis93806@cis.nctu.edu.tw

Jimmy J. M. Tan

Department of Computer Science  
National Chiao Tung University  
jmtan@cs.nctu.edu.tw

Lih-Hsing Hsu

Department of Computer Science and Information Engineering  
Providence University  
lhhsu@pu.edu.tw

## Abstract

In this article, we introduce 2RP-property in the augmented cube  $AQ_n$ : Let  $\{\mathbf{u}, \mathbf{v}, \mathbf{x}, \mathbf{y}\}$  be any four distinct vertices of  $AQ_n$ . Let  $l_1$  and  $l_2$  be two integers with  $l_1 \geq d_{AQ_n}(\mathbf{u}, \mathbf{v})$ ,  $l_2 \geq d_{AQ_n}(\mathbf{x}, \mathbf{y})$ , and  $l_1 + l_2 = 2^n - 2$ . Then there exist two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{u}$  and  $\mathbf{v}$  with  $l(P_1) = l_1$ , (2)  $P_2$  is a path joining  $\mathbf{x}$  and  $\mathbf{y}$  with  $l(P_2) = l_2$ , and (3)  $P_1 \cup P_2$  spans  $AQ_n$  except some special conditions.

*Keywords:* hamiltonian, augmented cubes.

Interconnection networks play an important role in parallel computing/communication systems. The graph embedding problem is a central issue in evaluating a network. The graph embedding problem asked if the guest graph is a subgraph of a host graph, and an important benefit of the graph embeddings is that we can apply existing algorithm for guest graphs to host graphs. This problem has attracted a burst of studies in recent years. Cycle networks and path networks are suitable for designing simple algorithms with low communication costs. The cycle embedding problem, which deals with all possible lengths of the cycles in a given graph, is investigated in a lot of inter-

## 1 Introduction

connection networks [6, 10, 12, 13]. The path embedding problem, which deals with all possible lengths of the paths between given two vertices in a given graph, is investigated in a lot of interconnection networks [3–5, 12–14].

In this article, a network is represented as a loopless undirected graph. For the graph definitions and notation, we follow [1]. Let  $G = (V, E)$  be a graph if  $V$  is a finite set and  $E$  is a subset of  $\{(a, b) \mid (a, b) \text{ is an unordered pair of } V\}$ . We say that  $V$  is the *vertex set* and  $E$  is the *edge set*. Two vertices  $u$  and  $v$  are *adjacent* if  $(u, v) \in E$ . We use  $Nbd_G(u)$  to denote the set  $\{v \mid (u, v) \in E(G)\}$ . The *degree* of a vertex  $u$  in  $G$ , denoted by  $\deg_G(u)$ , is  $|Nbd_G(u)|$ . We use  $\delta(G)$  to denote  $\min\{\deg_G(u) \mid u \in V(G)\}$ . A graph is *k-regular* if  $\deg_G(u) = k$  for every vertex  $u$  in  $G$ . A path is a sequence of adjacent vertices, written as  $\langle v_0, v_1, \dots, v_m \rangle$ , in which all the vertices  $v_0, v_1, \dots, v_m$  are distinct except that possibly  $v_0 = v_m$ . We also write the path  $\langle v_0, P, v_m \rangle$ , where  $P = \langle v_0, v_1, \dots, v_m \rangle$ . The *length* of a path  $P$ , denoted by  $l(P)$ , is the number of edges in  $P$ . Let  $u$  and  $v$  be two vertices of  $G$ . The *distance* between  $u$  and  $v$  denoted by  $d_G(u, v)$  is the length of the shortest path of  $G$  joining  $u$  and  $v$ . The *diameter* of a graph  $G$ , denoted by  $D(G)$ , is  $\max\{d_G(u, v) \mid u, v \in V(G)\}$ . A *cycle* is a path with at least three vertices such that the first vertex is the same as the last one. A *hamiltonian cycle* is a cycle of length  $V(G)$ . A *hamiltonian path* is a path of length  $V(G) - 1$ .

The hypercube  $Q_n$  is one of the most popular interconnection networks for parallel computer/communication system [11]. This is partly due to its attractive properties, such as regularity, recursive structure, vertex and edge symmetry, maximum connectivity, as well as effective routing and broadcasting algorithm. The augmented cube  $AQ_n$  is a variation of  $Q_n$ , proposed by Choudum and Sunitha [2], and not only retains some favorable prop-

erties of  $Q_n$  but also processes some embedding properties that  $Q_n$  does not [2, 7–9, 13]. For example,  $AQ_n$  contains cycles of all lengths from 3 to  $2^n$ , but  $Q_n$  contains only even cycles.

For the path embedding problem on the augmented cube, Ma et al. [13] proved that between any two distinct vertices  $\mathbf{x}$  and  $\mathbf{y}$  of  $AQ_n$ , there exists a path  $P_l(\mathbf{x}, \mathbf{y})$  of length  $l$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \leq l \leq 2^n - 1$ . Obviously, we expect that such a path  $P_l(\mathbf{x}, \mathbf{y})$  can be further extended by including the vertices not in  $P_l(\mathbf{x}, \mathbf{y})$  into a hamiltonian path from  $\mathbf{x}$  to a fixed vertex  $\mathbf{z}$  or a hamiltonian cycle. For this reason, we prove that for any three distinct vertices  $\mathbf{x}$ ,  $\mathbf{y}$  and  $\mathbf{z}$  of  $AQ_n$ , and for any  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \leq l \leq 2^n - 1 - d_{AQ_n}(\mathbf{y}, \mathbf{z})$  there exists a hamiltonian path  $R(\mathbf{x}, \mathbf{y}, \mathbf{z}; l)$  from  $\mathbf{x}$  to  $\mathbf{z}$  such that  $d_{R(\mathbf{x}, \mathbf{y}, \mathbf{z}; l)}(\mathbf{x}, \mathbf{y}) = l$ . As a corollary, we prove that for any two distinct vertices  $\mathbf{x}$  and  $\mathbf{y}$ , and for any  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \leq l \leq 2^{n-1}$ , there exists a hamiltonian cycle  $S(\mathbf{x}, \mathbf{y}; l)$  such that  $d_{S(\mathbf{x}, \mathbf{y}; l)}(\mathbf{x}, \mathbf{y}) = l$ .

In the following section, we introduce the definition and some properties of the augmented cubes. In Section 3, we introduce the 2RP-property for the augmented cube  $AQ_n$  and prove that  $AQ_n$  satisfies the 2RP-property if  $n \geq 2$ . We make some remarks to illustrate that some interesting properties of augmented cubes are consequences of 2RP-property in the final section.

## 2 Preliminaries

In this section, we introduce some properties of the augmented cubes. Assume that  $n \geq 1$  is an integer. The graph of the *n-dimensional augmented cube*, denoted by  $AQ_n$ , has  $2^n$  vertices, each labeled by an  $n$ -bit binary string  $V(AQ_n) = \{u_1u_2\dots u_n \mid u_i \in \{0, 1\}\}$ . For  $n = 1$ ,  $AQ_1$  is the graph  $K_2$  with vertex set  $\{0, 1\}$ . For  $n \geq 2$ ,  $AQ_n$  can be recursively constructed by two

copies of  $AQ_{n-1}$ , denoted by  $AQ_{n-1}^0$  and  $AQ_{n-1}^1$ , and by adding  $2^n$  edges between  $AQ_{n-1}^0$  and  $AQ_{n-1}^1$  as follows:

Let  $V(AQ_{n-1}^0) = \{0u_2u_3 \dots u_n \mid u_i = 0 \text{ or } 1 \text{ for } 2 \leq i \leq n\}$  and  $V(AQ_{n-1}^1) = \{1v_2v_3 \dots v_n \mid v_i = 0 \text{ or } 1 \text{ for } 2 \leq i \leq n\}$ . A vertex  $\mathbf{u} = 0u_2u_3 \dots u_n$  of  $AQ_{n-1}^0$  is adjacent to a vertex  $\mathbf{v} = 1v_2v_3 \dots v_n$  of  $AQ_{n-1}^1$  if and only if one of the following cases holds.

- (i)  $u_i = v_i$ , for  $2 \leq i \leq n$ . In this case,  $(\mathbf{u}, \mathbf{v})$  is called a *hypercube edge*. We set  $\mathbf{v} = \mathbf{u}^h$ .
- (ii)  $u_i = \bar{v}_i$ , for  $2 \leq i \leq n$ . In this case,  $(\mathbf{u}, \mathbf{v})$  is called a *complement edge*. We set  $\mathbf{v} = \mathbf{u}^c$ .

The augmented cubes  $AQ_1, AQ_2, AQ_3$  and  $AQ_4$  are illustrated in Figure 1. It is proved in [2] that  $AQ_n$  is a vertex transitive,  $(2n-1)$ -regular, and  $(2n-1)$ -connected graph with  $2^n$  vertices for any positive integer  $n$ . Let  $i$  be any index with  $1 \leq i \leq n$  and  $\mathbf{u} = u_1u_2u_3 \dots u_n$  be a vertex of  $AQ_n$ . We use  $\mathbf{u}^i$  to denote the vertex  $\mathbf{v} = v_1v_2v_3 \dots v_n$  such that  $u_j = v_j$  with  $1 \leq j \neq i \leq n$  and  $u_i = \bar{v}_i$ . Moreover, we use  $\mathbf{u}^{i*}$  to denote the vertex  $\mathbf{v} = v_1v_2v_3 \dots v_n$  such that  $u_j = v_i$  for  $j < i$  and  $u_j = \bar{v}_j$  for  $i \leq j \leq n$ . Obviously,  $\mathbf{u}^n = \mathbf{u}^{n*}$ ,  $\mathbf{u}^1 = \mathbf{u}^h$ ,  $\mathbf{u}^c = \mathbf{u}^{1*}$ , and  $Nbd_{AQ_n}(\mathbf{u}) = \{\mathbf{u}^i \mid 1 \leq i \leq n\} \cup \{\mathbf{u}^{i*} \mid 1 \leq i < n\}$ .

**Lemma 1.** Assume that  $n \geq 2$ . Then  $|Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{v})| \geq 2$  if  $(\mathbf{u}, \mathbf{v}) \in E(G)$ .

**Proof.** We prove this lemma by induction. Since  $AQ_2$  is isomorphic to the complete graph  $K_4$ , the lemma holds for  $n = 2$ . Assume the lemma holds for  $2 \leq k < n$ . Suppose that  $\{\mathbf{u}, \mathbf{v}\} \subset V(AQ_{n-1}^i)$  for some  $i \in \{0, 1\}$ . By induction,  $|Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{v})| \geq 2$ . Thus, consider the case that either  $\mathbf{v} = \mathbf{u}^h$  or  $\mathbf{v} = \mathbf{u}^c$ . Obviously,  $\{\mathbf{u}^{2*}, \mathbf{u}^c\} \subset Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{v})$  if  $\mathbf{v} = \mathbf{u}^h$ ; and




Figure 1: The augmented cubes  $AQ_1, AQ_2, AQ_3$  and  $AQ_4$ .

$\{\mathbf{u}^{2*}, \mathbf{u}^h\} \subset Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{v})$  if  $\mathbf{v} = \mathbf{u}^c$ . Then the statement holds.  $\square$

The following lemma can easily be obtained from the definition of  $AQ_n$ .

**Lemma 2.** Assume that  $n \geq 3$ . For any two different vertices  $\mathbf{u}$  and  $\mathbf{v}$  of  $AQ_n$ , there exists two other vertices  $\mathbf{x}$  and  $\mathbf{y}$  of  $AQ_n$  such that the subgraph of  $\{\mathbf{u}, \mathbf{v}, \mathbf{x}, \mathbf{y}\}$  containing a four cycle.

**Lemma 3.** [8] Let  $F$  be a subset of  $V(AQ_n)$ . Then there exists a hamiltonian path between any two vertices of  $V(AQ_n) - F$  if  $|F| \leq 2n - 4$  for  $n \geq 4$  and  $|F| \leq 1$  for  $n = 3$ .

**Lemma 4.** [2] Let  $\mathbf{u}$  and  $\mathbf{v}$  be any two vertices in  $AQ_n$  with  $n \geq 2$ . Suppose that both  $\mathbf{u}$  and  $\mathbf{v}$  are in  $AQ_{n-1}^i$  for  $i = 0, 1$ . Then  $d_{AQ_n}(\mathbf{u}, \mathbf{v}) = d_{AQ_{n-1}^i}(\mathbf{u}, \mathbf{v})$ . Suppose that  $\mathbf{u}$  is a vertex in  $AQ_{n-1}^i$  and  $\mathbf{v}$  is a vertex in  $AQ_{n-1}^{1-i}$ . Then there exist two shortest paths  $P_1$  and  $P_2$  of  $AQ_n$  joining  $\mathbf{u}$  to  $\mathbf{v}$  such that  $(V(P_1) - \{\mathbf{v}\}) \subset V(AQ_{n-1}^i)$  and  $(V(P_2) - \{\mathbf{u}\}) \subset V(AQ_{n-1}^{1-i})$ .

With Lemma 4, we have the following corollary.

**Corollary 5.** Assume that  $n \geq 3$ . Let  $\mathbf{x}$  and  $\mathbf{y}$  be two vertices of  $AQ_n$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \geq 2$ . Then, there are

two vertices  $\mathbf{p}$  and  $\mathbf{q}$  in  $Nbd_{AQ_n}(\mathbf{x})$  with  $d_{AQ_n}(\mathbf{p}, \mathbf{y}) = d_{AQ_n}(\mathbf{q}, \mathbf{y}) = d_{AQ_n}(\mathbf{x}, \mathbf{y}) - 1$ .

**Lemma 6.** [8] *Let  $\{\mathbf{u}, \mathbf{v}, \mathbf{x}, \mathbf{y}\}$  be any four distinct vertices of  $AQ_n$  with  $n \geq 2$ . Then there exist two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{u}$  and  $\mathbf{v}$ , (2)  $P_2$  is a path joining  $\mathbf{x}$  and  $\mathbf{y}$ , and (3)  $P_1 \cup P_2$  spans  $AQ_n$ .*

We refer to Lemma 6 as 2P-property of the augmented cube. This property is used for many applications of the augmented cubes [7, 8]. Obviously,  $l(P_1) \geq d_{AQ_n}(\mathbf{u}, \mathbf{v})$  and  $l(P_2) \geq d_{AQ_n}(\mathbf{x}, \mathbf{y})$ , and  $l(P_1) + l(P_2) = 2^n - 2$ . We expect that  $l(P_1)$ , hence,  $l(P_2)$  can be an arbitrarily integer with the above constraint. However, such expectation is almost true. Let us consider  $AQ_3$ . Suppose that  $\mathbf{u} = 001$ ,  $\mathbf{v} = 110$ ,  $\mathbf{x} = 101$ , and  $\mathbf{y} = 010$ . Thus,  $d_{AQ_3}(\mathbf{u}, \mathbf{v}) = 1$  and  $d_{AQ_3}(\mathbf{x}, \mathbf{y}) = 1$ . We can find  $P_1$  and  $P_2$  with  $l(P_1) \in \{1, 3, 5\}$ . Note that  $\{\mathbf{x}, \mathbf{y}\} = Nbd_{AQ_3}(\mathbf{u}) \cap Nbd_{AQ_3}(\mathbf{v})$ . We can not find  $P_1$  with  $l(P_1) = 2$ . Again,  $\{\mathbf{u}, \mathbf{v}\} = Nbd_{AQ_3}(\mathbf{x}) \cap Nbd_{AQ_3}(\mathbf{y})$ . We can not find  $P_2$  with  $l(P_2) = 2$ . Hence, we cannot find  $P_1$  with  $l(P_1) = 4$ . Similarly, we consider  $AQ_4$ . Suppose that  $\mathbf{u} = 0000$ ,  $\mathbf{v} = 1001$ ,  $\mathbf{x} = 0001$  and  $\mathbf{y} = 1000$ . Thus,  $d_{AQ_4}(\mathbf{u}, \mathbf{v}) = 2$  and  $d_{AQ_4}(\mathbf{x}, \mathbf{y}) = 2$ . We can find  $P_1$  and  $P_2$  with  $l(P_1) \in \{3, 4, \dots, 11\}$ . Note that  $\{\mathbf{x}, \mathbf{y}\} = Nbd_{AQ_4}(\mathbf{u}) \cap Nbd_{AQ_4}(\mathbf{v})$ . We can not find  $P_1$  with  $l(P_1) = 2$ . Again,  $\{\mathbf{u}, \mathbf{v}\} = Nbd_{AQ_4}(\mathbf{x}) \cap Nbd_{AQ_4}(\mathbf{y})$ . We can not find  $P_2$  with  $l(P_2) = 2$ .

### 3 The 2RP-property of the augmented cubes

In this section, we introduce the 2RP-property for the augmented cube  $AQ_n$  and prove that  $AQ_n$  satisfies the 2RP-property if  $n \geq 2$ . First, we propose the 2RP-property of  $AQ_n$  with  $n \geq 2$ : *Let  $\{\mathbf{u}, \mathbf{v}, \mathbf{x}, \mathbf{y}\}$  be any*

*four distinct vertices of  $AQ_n$ . Let  $l_1$  and  $l_2$  be two integers with  $l_1 \geq d_{AQ_n}(\mathbf{u}, \mathbf{v})$ ,  $l_2 \geq d_{AQ_n}(\mathbf{x}, \mathbf{y})$ , and  $l_1 + l_2 = 2^n - 2$ . Then there exist two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{u}$  and  $\mathbf{v}$  with  $l(P_1) = l_1$ , (2)  $P_2$  is a path joining  $\mathbf{x}$  and  $\mathbf{y}$  with  $l(P_2) = l_2$ , and (3)  $P_1 \cup P_2$  spans  $AQ_n$  except for the following cases: (a)  $l_1 = 2$  with  $d_{AQ_n}(\mathbf{u}, \mathbf{v}) = 1$  such that  $\{\mathbf{x}, \mathbf{y}\} = Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{v})$ ; (b)  $l_2 = 2$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$  such that  $\{\mathbf{u}, \mathbf{v}\} = Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ ; (c)  $l_1 = 2$  with  $d_{AQ_n}(\mathbf{u}, \mathbf{v}) = 2$  such that  $\{\mathbf{x}, \mathbf{y}\} = Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{v})$ ; and (d)  $l_2 = 2$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 2$  such that  $\{\mathbf{u}, \mathbf{v}\} = Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ .*

**Theorem 7.** *Assume that  $n$  is a positive integer with  $n \geq 2$ . Then  $AQ_n$  satisfies 2RP-property.*

**Proof.** We prove this theorem by induction. By brute force, we check the theorem holds for  $n = 2, 3, 4$ . Assume the theorem holds for any  $AQ_k$  with  $4 \leq k < n$ . Without loss of generality, we can assume that  $l_1 \geq l_2$ . Thus,  $l_2 \leq 2^{n-1} - 1$ . By the symmetric property of  $AQ_n$ , we can assume that at least one of  $\mathbf{u}$  and  $\mathbf{v}$ , say  $\mathbf{u}$ , is in  $V(AQ_{n-1}^0)$ . Thus, we have the following cases:

**Case 1:**  $\mathbf{v} \in V(AQ_{n-1}^0)$  and  $\{\mathbf{x}, \mathbf{y}\} \subset V(AQ_{n-1}^1)$ .

**Subcase 1.1:**  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \leq l_2 \leq 2^{n-1} - 3$  except that (1)  $l_2 = 2^{n-1} - 4$  and (2)  $l_2 = 2$  if  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$  or 2 with  $\{\mathbf{u}, \mathbf{v}\} \neq Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . See Figure 2(a) for an illustration. By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Since  $l(R) = 2^{n-1} - 1$ , we can write  $R$  as  $\langle \mathbf{u}, R_1, \mathbf{p}, \mathbf{q}, R_2, \mathbf{v} \rangle$  for some vertices  $\mathbf{p}$  and  $\mathbf{q}$  such that  $\{\mathbf{p}^h, \mathbf{q}^h\} \cap \{\mathbf{x}, \mathbf{y}\} = \emptyset$ . By induction, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{p}^h$  to  $\mathbf{q}^h$  with  $l(S_1) = 2^{n-1} - l_2 - 2$ , (2)  $S_2$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$  with  $l(S_2) = l_2$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set

$P_1$  as  $\langle u, R_1, p, p^h, S_1, q^h, q, R_2, v \rangle$  and set  $P_2$  as  $S_2$ . Obviously,  $P_1$  and  $P_2$  are the required paths.





Figure 2: Subcase 1.1 and Subcase 1.3.

**Subcase 1.2:**  $l_2 = 2$  if  $d_{AQ_n}(x, y) = 1$  or  $2$  with  $\{u, v\} \neq Nbd_{AQ_n}(x) \cap Nbd_{AQ_n}(y)$ . Obviously, there exists a path  $P_2$  of length 2 in  $AQ_n - \{u, v\}$  joining  $x$  to  $y$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - V(P_2)$  joining  $u$  to  $v$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 1.3:**  $l_2 = 2^{n-1} - 4$ . See Figure 2(b) for an illustration. Obviously, there exists a vertex  $p$  in  $V(AQ_{n-1}^1) - \{x, y, u^h, v^h\}$ , a vertex  $q$  in  $Nbd_{AQ_{n-1}^1}(p) - \{x, y\}$ ,

and a vertex  $r$  in  $Nbd_{AQ_{n-1}^1}(q) - \{x, y, p\}$ . Suppose that  $r^h \notin \{u, v\}$ . By induction, there exist two disjoint paths  $Q_1$  and  $Q_2$  such that (1)  $Q_1$  is a path joining  $u$  to  $p^h$ , (2)  $Q_2$  is a path joining  $r^h$  to  $v$ , and (3)  $Q_1 \cup Q_2$  spans  $AQ_{n-1}^0$ . By Lemma 3, there exists a hamiltonian path  $P_2$  of  $AQ_{n-1}^1 - \{p, q, r\}$  joining  $x$  to  $y$ . We set  $P_1$  as  $\langle u, Q_1, p^h, p, q, r, r^h, Q_2, v \rangle$ . Suppose that  $r^h \in \{u, v\}$ . Without loss of generality, we assume that  $r^h = v$ . By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0 - \{v\}$  joining  $u$  to  $p^h$ . We set  $P_1$  as  $\langle u, R, p^h, p, q, r, r^h = v \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 1.4:**  $l_2 = 2^{n-1} - 2$ . Obviously, there exist a vertex  $p \in V(AQ_{n-1}^1) - \{x, y, u^h, u^c, v^h, v^c\}$ . By Lemma 6, there exist two disjoint paths  $Q_1$  and  $Q_2$  such that (1)  $Q_1$  is a path joining  $u$  and  $p^h$ , (2)  $Q_2$  is a path joining  $p^c$  and  $v$ , and (3)  $Q_1 \cup Q_2$  spans  $AQ_{n-1}^0$ . By Lemma 3, there exists a hamiltonian path  $P_2$  of  $AQ_{n-1}^0 - \{p\}$  joining  $x$  to  $y$ . We set  $P_1$  as  $\langle u, Q_1, p^h, p, p^c, Q_2, v \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 1.5:**  $l_2 = 2^{n-1} - 1$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_{n-1}^0$  joining  $u$  and  $v$  and there exists a hamiltonian path  $P_2$  of  $AQ_{n-1}^1$  joining  $x$  to  $y$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Case 2:**  $v \in V(AQ_{n-1}^0)$  and exactly one of  $x$  and  $y$  is in  $V(AQ_{n-1}^0)$ . Without loss of generality, we assume that  $x \in V(AQ_{n-1}^0)$ .

**Subcase 2.1:**  $l_2 = 1$ . Obviously,  $d_{AQ_n}(x, y) = 1$ . We set  $P_2$  as  $\langle x, y \rangle$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - \{x, y\}$  joining  $u$  to  $v$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 2.2:**  $l_2 = 2$  if  $d_{AQ_n}(x, y) = 1$  or  $2$  with  $\{u, v\} \neq Nbd_{AQ_n}(x) \cap Nbd_{AQ_n}(y)$ . The proof is the same to Subcase 1.2.

**Subcase 2.3:**  $l_2 = 3$ .

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$ . There exists a vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{x}) - \{\mathbf{u}, \mathbf{v}\}$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - \{\mathbf{x}, \mathbf{y}, \mathbf{p}, \mathbf{p}^h\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, \mathbf{p}^h, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 2$ . By Lemma 4, there exists a path  $\langle \mathbf{x}, \mathbf{p}, \mathbf{y} \rangle$  from  $\mathbf{x}$  to  $\mathbf{y}$  such that  $\mathbf{p} \in V(AQ_{n-1}^1)$ . By Lemma 1, there exists a vertex  $\mathbf{q} \in Nbd_{AQ_{n-1}^1}(\mathbf{p}) \cap Nbd_{AQ_{n-1}^1}(\mathbf{y})$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - \{\mathbf{x}, \mathbf{y}, \mathbf{p}, \mathbf{q}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, \mathbf{q}, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 3$ . By Lemma 4, there exists a path  $P_2$  from  $\mathbf{x}$  to  $\mathbf{y}$  such that  $(V(P_2) - \{\mathbf{x}\}) \subset V(AQ_{n-1}^1)$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - V(P_2)$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 2.4:**  $4 \leq l_2 \leq 2^{n-1} - 2$  except that  $l_2 = 2^{n-1} - 3$ .

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$  or  $2$ . We first claim that there exists a vertex  $\mathbf{p}$  in  $Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . Assume that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$ . Obviously, either  $\mathbf{y} = \mathbf{x}^h$  or  $\mathbf{y} = \mathbf{x}^c$ . We set  $\mathbf{p} = \mathbf{x}^c$  if  $\mathbf{y} = \mathbf{x}^h$ ; and we set  $\mathbf{p} = \mathbf{x}^h$  if  $\mathbf{y} = \mathbf{x}^c$ . Assume that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 2$ . See Figure 3 for an illustration. By Lemma 4, there exists a path  $\langle \mathbf{x}, \mathbf{p}, \mathbf{y} \rangle$  from  $\mathbf{x}$  to  $\mathbf{y}$  such that  $\mathbf{p} \in V(AQ_{n-1}^1)$ . Obviously,  $\mathbf{p}$  satisfies our claim. By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0 - \{\mathbf{x}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Since  $l(R) = 2^{n-1} - 3$ , we can write  $R$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{t}, R_2, \mathbf{v} \rangle$  such that  $\{\mathbf{s}^h, \mathbf{t}^h\} \cap \{\mathbf{p}, \mathbf{y}\} = \emptyset$ . By induction, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{s}^h$  to  $\mathbf{t}^h$  with  $l(S_1) = 2^{n-1} - 1 - l_2$ , (2)  $S_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{y}$  with  $l(S_2) = l_2 - 1$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^h, S_1, \mathbf{t}^h, \mathbf{t}, R_2, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, S_2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required

paths.




Figure 3: Subcase 2.4.

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \geq 3$ . By Lemma 4, there exists a vertex  $\mathbf{p}$  in  $V(AQ_{n-1}^1)$  such that  $d_{AQ_n}(\mathbf{p}, \mathbf{y}) = d_{AQ_n}(\mathbf{x}, \mathbf{y}) - 1$ . By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0 - \{\mathbf{x}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . We can write  $R$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{t}, R_2, \mathbf{v} \rangle$  such that  $\{\mathbf{s}^h, \mathbf{t}^h\} \cap \{\mathbf{p}, \mathbf{y}\} = \emptyset$ . By induction, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{s}^h$  to  $\mathbf{t}^h$  with  $l(S_1) = 2^{n-1} - 1 - l_2$ , (2)  $S_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{y}$  with  $l(S_2) = l_2 - 1$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^h, S_1, \mathbf{t}^h, \mathbf{t}, R_2, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, S_2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 2.5:**  $l_2 = 2^{n-1} - 3$  or  $l_2 = 2^{n-1} - 1$ . Let  $k = 3$  if  $l_2 = 2^{n-1} - 3$  and  $k = 1$  if  $l_2 = 2^{n-1} - 1$ . There exists a vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{x}) - \{\mathbf{u}, \mathbf{v}, \mathbf{y}^n\}$ . By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0 - \{\mathbf{x}, \mathbf{p}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . We can write  $R$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{t}, R_2, \mathbf{v} \rangle$  such that  $\{\mathbf{s}, \mathbf{t}\} \cap \{\mathbf{p}, \mathbf{y}^n\} = \emptyset$ . By induction, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{s}^n$  to  $\mathbf{t}^n$  with  $l(S_1) = k$ , (2)  $S_2$  is a path joining  $\mathbf{p}^n$  to  $\mathbf{y}$  with  $l(S_2) = 2^{n-1} - k - 2$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^n, S_1, \mathbf{t}^n, \mathbf{t}, R_2, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, \mathbf{p}^n, S_2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the re-

quired paths.

**Case 3:**  $\{v, x, y\} \subset V(Q_{n-1}^0)$ .

**Subcase 3.1:**  $l_2 = 1$ . The proof is the same as Subcase 2.1.

**Subcase 3.2:**  $l_2 = 2$  if  $d_{AQ_n}(x, y) = 1$  or 2 with  $\{u, v\} \neq Nbd_{AQ_n}(x) \cap Nbd_{AQ_n}(y)$ . The proof is the same as Subcase 1.2.

**Subcase 3.3:**  $d_{AQ_n}(x, y) \leq l_2 \leq 2^{n-2} - 1$ . See Figure 4(a) for an illustration. By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $u$  to  $v$  with  $l(R_1) = 2^{n-1} - l_2 - 2$ , (2)  $R_2$  is a path joining  $x$  to  $y$  with  $l(R_2) = l_2$ , (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . We can write  $R_1$  as  $\langle u, R_3, p, q, R_4, v \rangle$ . By Lemma 3, there exists a hamiltonian path  $S$  of  $AQ_{n-1}^1$  joining  $p^h$  to  $q^h$ . We set  $P_1$  as  $\langle u, R_3, p, p^h, S, q^h, q, R_4, v \rangle$  and  $P_2$  as  $R_2$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 3.4:**  $2^{n-2} + 1 \leq l_2 \leq 2^{n-1} - 1$  except that  $l_2 = 2^{n-2} + 2$ . See Figure 4(b) for an illustration. By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $u$  to  $v$  with  $l(R_1) = 2^{n-2} - 1$ , (2)  $R_2$  is a path joining  $x$  to  $y$  with  $l(R_2) = 2^{n-2} - 1$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . We can write  $R_1$  as  $\langle u, R_3, p, q, R_4, v \rangle$  and write  $R_2$  as  $\langle x, R_5, s, t, R_6, y \rangle$ . By induction, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $p^h$  to  $q^h$  with  $l(S_1) = 2^{n-1} - l_2 + 2^{n-2} - 2$ , (2)  $S_2$  is a path joining  $s^h$  to  $t^h$  with  $l(S_2) = l_2 - 2^{n-2}$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set  $P_1$  as  $\langle u, R_3, p, p^h, S_1, q^h, q, R_4, v \rangle$  and  $P_2$  as  $\langle x, R_5, s, s^h, S_2, t^h, t, R_6, y \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 3.5:**  $l_2 = 2^{n-2}$  or  $2^{n-2} + 2$ . Let  $k = 0$  if  $l_2 = 2^{n-2}$  and  $k = 2$  if  $l_2 = 2^{n-2} + 2$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $u$  to  $v$  with  $l(R_1) = 2^{n-2} - k$ , (2)  $R_2$  is a path joining  $x$  to  $y$  with  $l(R_2) = 2^{n-2} + k - 2$ , and (3)  $R_1 \cup R_2$

spans  $AQ_{n-1}^0$ . We can write  $R_1$  as  $\langle u, R_3, p, q, R_4, v \rangle$  and write  $R_2$  as  $\langle x, R_5, s, t, R_6, y \rangle$ . By Lemma 3, there exists a hamiltonian path  $S$  of  $AQ_{n-1}^1 - \{s^n, t^n\}$  joining  $p^n$  to  $q^n$ . We set  $P_1$  as  $\langle u, R_3, p, p^n, S, q^n, q, R_4, v \rangle$  and  $P_2$  as  $\langle x, R_5, s, s^n, t^n, t, R_6, y \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.





Figure 4: Subcase 3.3 and Subcase 3.4.


**Case 4:**  $\{x, v, y\} \subset V(AQ_{n-1}^1)$ .

**Subcase 4.1:**  $d_{AQ_n}(x, y) \leq l_2 \leq 2^{n-1} - 3$  except that (1)  $l_2 = 2^{n-1} - 4$  and (2)  $l_2 = 2$  if  $d_{AQ_n}(x, y) = 1$  or 2 with  $\{u, v\} \neq Nbd_{AQ_n}(x) \cap Nbd_{AQ_n}(y)$ . See Figure 5(a) for an illustration. Obviously, there exists a vertex  $p$  in  $Nbd_{AQ_{n-1}^1}(v) - \{x, y, u^h\}$ . By induction, there exist two

disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{p}$  to  $\mathbf{v}$  with  $l(S_1) = l_1 - 2^{n-1}$ , (2)  $S_2$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$  with  $l(S_2) = l_2$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0$  joining  $\mathbf{u}$  and  $\mathbf{p}^h$ . We set  $P_1$  as  $\langle \mathbf{u}, R, \mathbf{p}^h, \mathbf{p}, S_1, \mathbf{v} \rangle$  and we set  $P_2$  as  $S_2$ . Obviously,  $P_1$  and  $P_2$  are the required paths.



(a)



(b)

Figure 5: Subcase 4.1 and Subcase 4.5.

**Subcase 4.2:**  $l_2 = 2$  if  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$  or  $2$  with  $\{\mathbf{u}, \mathbf{v}\} \neq Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . The proof is the same to Subcase 1.2.

**Subcase 4.3:**  $l_2 = 2^{n-1} - 4$ . Obviously, there exists a

vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^1}(\mathbf{v}) - \{\mathbf{x}, \mathbf{y}\}$ , and there exists a vertex  $\mathbf{q}$  in  $Nbd_{AQ_{n-1}^1}(\mathbf{p}) - \{\mathbf{x}, \mathbf{y}, \mathbf{v}, \mathbf{u}^h\}$ . By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0$  joining  $\mathbf{u}$  to  $\mathbf{q}^h$ , and there exists a hamiltonian path  $P_2$  of  $AQ_{n-1}^1 - \{\mathbf{v}, \mathbf{p}, \mathbf{q}\}$  joining  $\mathbf{x}$  to  $\mathbf{y}$ . We set  $P_1$  as  $\langle \mathbf{u}, R, \mathbf{q}^h, \mathbf{q}, \mathbf{p}, \mathbf{v} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 4.4:**  $l_2 = 2^{n-1} - 2$ . Let  $\mathbf{v}'$  be an element in  $\{\mathbf{v}^h, \mathbf{v}^c\} - \{\mathbf{u}\}$ . By Lemma 3, there exists a hamiltonian path  $R$  of  $AQ_{n-1}^0$  joining  $\mathbf{u}$  to  $\mathbf{v}'$ , and there exists a hamiltonian path  $P_2$  of  $AQ_{n-1}^1 - \{\mathbf{v}\}$  joining  $\mathbf{x}$  to  $\mathbf{y}$ . We set  $P_1$  as  $\langle \mathbf{u}, R, \mathbf{v}', \mathbf{v} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 4.5:**  $l_2 = 2^{n-1} - 1$ . See Figure 5(b) for an illustration. Obviously, there exists a vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^1}(\mathbf{v}) - \{\mathbf{x}, \mathbf{y}\}$ . By induction, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{p}$  to  $\mathbf{v}$  with  $l(S_1) = 1$ , (2)  $S_2$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$  with  $l(S_2) = 2^{n-1} - 3$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . Obviously, we can write  $S_2$  as  $\langle \mathbf{x}, S_2^1, \mathbf{r}, \mathbf{s}, S_2^2, \mathbf{y} \rangle$  for some vertex  $\mathbf{r}$  and  $\mathbf{s}$  such that  $\mathbf{u} \notin \{\mathbf{r}^h, \mathbf{s}^h\}$ . Again by induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{p}^h$  with  $l(R_1) = 2^{n-1} - 3$ , (2)  $R_2$  is a path joining  $\mathbf{r}^h$  to  $\mathbf{s}^h$  with  $l(R_2) = 1$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{p}^h, \mathbf{p}, \mathbf{v} \rangle$  and set  $P_2$  as  $\langle \mathbf{x}, S_2^1, \mathbf{r}, \mathbf{r}^h, \mathbf{s}^h, \mathbf{s}, S_2^2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Case 5:**  $\mathbf{v} \in V(AQ_{n-1}^1)$  and  $|\{x, y\} \cap V(AQ_{n-1}^0)| = 1$ . Without loss of generality, we assume that  $\mathbf{x} \in V(AQ_{n-1}^0)$ .

**Subcase 5.1:**  $l_2 = 1$ . The proof is the same to Subcase 2.1.

**Subcase 5.2:**  $l_2 = 2$  if  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$  or  $2$  with  $\{\mathbf{u}, \mathbf{v}\} \neq Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . The proof is the same to Subcase 1.2.

**Subcase 5.3:**  $l_2 = 3$ .



Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$ . Obviously, there exists a vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{x}) - \{\mathbf{u}, \mathbf{v}^h\}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, \mathbf{p}^h, \mathbf{y} \rangle$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - V(P_2)$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 2$ . Assume that  $\{\mathbf{u}, \mathbf{v}\} = Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . Thus, we have either  $\mathbf{v} = \mathbf{x}^h$  or  $\mathbf{v} = \mathbf{x}^c$ . Moreover,  $\mathbf{u} = \mathbf{x}^\alpha$ , and  $\mathbf{y} = \mathbf{v}^\alpha$  for some  $\alpha \in \{i \mid 2 \leq i \leq n\} \cup \{i^* \mid 2 \leq i \leq n-1\}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{x}^{h*}, (\mathbf{x}^{h*})^\alpha, ((\mathbf{x}^h)^\alpha) = \mathbf{y} \rangle$  in the case of  $\mathbf{v} = \mathbf{x}^h$ . Otherwise, we set  $P_2$  as  $\langle \mathbf{x}, \mathbf{x}^h, (\mathbf{x}^h)^\alpha, ((\mathbf{x}^{h*})^\alpha) = \mathbf{y} \rangle$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - V(P_2)$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Obviously,  $P_1$  and  $P_2$  are the required paths. Now, assume that  $\{\mathbf{u}, \mathbf{v}\} \neq Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . By Lemma 1, there exists a vertex  $\mathbf{p}$  in  $(Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})) - \{\mathbf{u}, \mathbf{v}\}$ . Without loss of generality, we may assume that  $\mathbf{p}$  is in  $AQ_{n-1}^0$ . By Lemma 1, there exists a vertex  $\mathbf{q}$  in  $(Nbd_{AQ_{n-1}^0}(\mathbf{p}) \cap Nbd_{AQ_{n-1}^0}(\mathbf{x})) - \{\mathbf{u}\}$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - \{\mathbf{x}, \mathbf{q}, \mathbf{p}, \mathbf{y}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{q}, \mathbf{p}, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 3$ . By Lemma 4, there are two shortest paths  $R_1$  and  $R_2$  of  $AQ_n$  joining  $\mathbf{x}$  to  $\mathbf{y}$  such that  $R_1$  can be written as  $\langle \mathbf{x}, \mathbf{r}_1, \mathbf{r}_2, \mathbf{y} \rangle$  with  $\{\mathbf{r}_1, \mathbf{r}_2\} \subset V(AQ_{n-1}^0)$  and  $R_2$  can be written as  $\langle \mathbf{x}, \mathbf{s}_1, \mathbf{s}_2, \mathbf{y} \rangle$  with  $\{\mathbf{s}_1, \mathbf{s}_2\} \subset V(AQ_{n-1}^1)$ . Suppose that  $\mathbf{u} \neq \mathbf{r}_2$  or  $\mathbf{v} \neq \mathbf{s}_1$ . Without loss of generality, we assume that  $\mathbf{u} \neq \mathbf{r}_2$ . By Corollary 5, there exists a vertex  $\mathbf{t} \in Nbd_{AQ_{n-1}^0}(\mathbf{x}) \cap Nbd_{AQ_{n-1}^0}(\mathbf{r}_2) - \{\mathbf{u}\}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{t}, \mathbf{r}_2, \mathbf{y} \rangle$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - V(P_2)$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Obviously,  $P_1$  and  $P_2$  are the required paths. Thus, we consider  $\mathbf{u} = \mathbf{r}_2$  and  $\mathbf{v} = \mathbf{s}_1$ . By Corollary 5, there exists a vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{x}) \cap Nbd_{AQ_{n-1}^0}(\mathbf{u})$ . Obviously,

$d_{AQ_n}(\mathbf{p}, \mathbf{y}) = 2$ . By Lemma 4, there exists a vertex  $\mathbf{q}$  in  $V(AQ_{n-1}^1) \cap Nbd_{AQ_n}(\mathbf{p}) \cap Nbd_{AQ_n}(\mathbf{y})$ . Since  $d_{AQ_n}(\mathbf{q}, \mathbf{y}) = 1$  and  $d_{AQ_n}(\mathbf{v}, \mathbf{y}) = 2$ ,  $\mathbf{q} \neq \mathbf{v}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, \mathbf{q}, \mathbf{y} \rangle$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - V(P_2)$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 5.4:**  $4 \leq l_2 \leq 2^{n-1} - 1$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 1$ .

Suppose that  $l_2 = 4$ . See Figure 6(a) for an illustration. Obviously, there exists a vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{x}) - \{\mathbf{u}, \mathbf{v}^h\}$ . By Lemma 1, there exists a vertex  $\mathbf{q}$  in  $(Nbd_{AQ_{n-1}^0}(\mathbf{x}) \cap Nbd_{AQ_{n-1}^0}(\mathbf{p})) - \{\mathbf{u}\}$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - \{\mathbf{x}, \mathbf{y}, \mathbf{p}, \mathbf{p}^h, \mathbf{q}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{q}, \mathbf{p}, \mathbf{p}^h, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $5 \leq l_2 \leq 2^{n-1} - 1$  except that  $l_2 = 2^{n-1} - 2$ . See Figure 6(b) for an illustration. Obviously, there exist a vertex  $\mathbf{p}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{x}) - \{\mathbf{u}, \mathbf{v}^h, \mathbf{y}^h\}$  and a vertex  $\mathbf{s}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{u}) - \{\mathbf{x}, \mathbf{p}, \mathbf{v}^h, \mathbf{y}^h\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{s}$  with  $l(R_1) = 2^{n-1} - 2 - l_2$ , (2)  $R_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{x}$  with  $l(R_2) = l_2 - 2$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . By Lemma 3, there exists a hamiltonian path  $S$  of  $AQ_{n-1}^1 - \{\mathbf{y}, \mathbf{p}^h\}$  joining  $\mathbf{s}^h$  to  $\mathbf{v}$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^h, S, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, R_2, \mathbf{p}, \mathbf{p}^h, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $l_2 = 2^{n-1} - 2$ . See Figure 6(c) for an illustration. Let  $\mathbf{s}$  and  $\mathbf{p}$  be two vertices in  $V(AQ_{n-1}^0) - \{\mathbf{u}, \mathbf{x}, \mathbf{v}^h, \mathbf{y}^h\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{s}$  with  $l(R_1) = 2^{n-2}$ , (2)  $R_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{x}$  with  $l(R_2) = 2^{n-2} - 2$ , (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . Similarly, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{s}^h$  to  $\mathbf{v}$  with  $l(S_1) = 2^{n-2} - 1$ , (2)  $S_2$  is a path joining  $\mathbf{p}^h$  to  $\mathbf{y}$  with  $l(S_2) = 2^{n-2} - 1$ , and (3)




Figure 6: Subcase 5.4.

$S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set  $P_1$  as  $\langle u, R_1, s, s^h, S_1, v \rangle$  and  $P_2$  as  $\langle x, R_2, p, p^h, S_2, y \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 5.5:**  $4 \leq l_2 \leq 2^{n-1} - 1$  except  $l_2 = 2^{n-1} - 3$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \geq 2$ .

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 2$  with  $\{\mathbf{u}, \mathbf{v}\} = Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . See Figure 7(a) for an illustration. Thus, we have either  $\mathbf{v} = \mathbf{x}^h$  or  $\mathbf{v} = \mathbf{x}^c$ . Moreover,  $\mathbf{u} = \mathbf{x}^\alpha$  and  $\mathbf{y} = (\mathbf{x}^h)^\alpha$  for some  $\alpha \in \{i \mid 2 \leq i \leq n\} \cup \{i^* \mid 2 \leq i \leq n-1\}$ . Obviously, there exists a vertex  $\mathbf{t}$  in  $Nbd_{AQ_{n-1}^1}(\mathbf{v}) - \{\mathbf{x}^h, \mathbf{y}, \mathbf{x}^c, \mathbf{u}^h\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{t}$  to  $\mathbf{v}$  with  $l(R_1) = 2^{n-1} - 1 - l_2$ , (2)  $R_2$  is a path joining  $\mathbf{x}^c$  to  $\mathbf{y}$  with  $l(R_2) = l_2 - 1$  in the case of  $\mathbf{v} = \mathbf{x}^h$ ; otherwise  $R_2$  is a path joining  $\mathbf{x}^h$  to  $\mathbf{y}$  with  $l(R_2) = l_2 - 1$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^1$ . By Lemma 3, there exists a hamiltonian path  $S$  of  $AQ_{n-1}^0 - \{\mathbf{x}\}$  joining  $\mathbf{t}^h$  to  $\mathbf{u}$ . We set  $P_1$  as  $\langle \mathbf{u}, S, \mathbf{t}^h, \mathbf{t}, R_1, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, \mathbf{x}^c, R_2, \mathbf{y} \rangle$  in the case of  $\mathbf{v} = \mathbf{x}^h$ ; otherwise, we set  $P_2$  as  $\langle \mathbf{x}, \mathbf{x}^h, R_2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 2$  with  $\{\mathbf{u}, \mathbf{v}\} \neq Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . See Figure 7(b) for an illustration. Then, there exists a vertex  $\mathbf{p}$  in  $(Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})) - \{\mathbf{u}, \mathbf{v}\}$ . Without loss of generality, we may assume that  $\mathbf{p} \in V(AQ_{n-1}^1)$ . Obviously, there exists a vertex  $\mathbf{t}$  in  $Nbd_{AQ_{n-1}^1}(\mathbf{v}) - \{\mathbf{y}, \mathbf{p}, \mathbf{u}^h, \mathbf{x}^h\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{t}$  to  $\mathbf{v}$  with  $l(R_1) = 2^{n-1} - 1 - l_2$ , (2)  $R_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{y}$  with  $l(R_2) = l_2 - 1$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^1$ . By Lemma 3, there exists a hamiltonian path  $S$  of  $AQ_{n-1}^0 - \{\mathbf{x}\}$  joining  $\mathbf{t}^h$  to  $\mathbf{u}$ . We set  $P_1$  as  $\langle \mathbf{u}, S, \mathbf{t}^h, \mathbf{t}, R_1, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, \mathbf{p}, R_2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Suppose that  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = k \geq 3$ . By




Figure 7: Subcase 5.5.

Lemma 4, there are two shortest paths  $S_1$  and  $S_2$  of  $AQ_n$  joining  $\mathbf{x}$  to  $\mathbf{y}$  such that  $S_1$  can be written as  $\langle \mathbf{x} = \mathbf{r}_0, \mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_{k-1}, \mathbf{y} \rangle$  with  $(V(S_1) - \{\mathbf{y}\}) \subset V(AQ_{n-1}^0)$  and  $S_2$  can be written as  $\langle \mathbf{x}, \mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_{k-1}, \mathbf{y} \rangle$  with  $(V(S_2) - \{\mathbf{x}\}) \subset V(AQ_{n-1}^1)$ . Suppose that  $\mathbf{u} \neq \mathbf{r}_{k-1}$ . See Figure 7(c) for an illustration. We set  $\mathbf{p} = \mathbf{r}_{k-1}$ . Again, there exists a vertex  $\mathbf{s}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{u}) - \{\mathbf{x}, \mathbf{p}, \mathbf{y}^h, \mathbf{v}^h\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{s}$  with  $l(R_1) = 2^{n-1} - 1 - l_2$ , (2)  $R_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{x}$  with  $l(R_2) = l_2 - 1$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . By Lemma 3, there exists a hamiltonian path  $S$  of  $AQ_{n-1}^1 - \{\mathbf{y}\}$  joining  $\mathbf{s}^h$  to  $\mathbf{v}$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^h, S, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, R_2, \mathbf{p}, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Now we assume that  $\mathbf{r}_{k-1} = \mathbf{u}$  and  $\mathbf{s}_1 = \mathbf{v}$ . See Figure 7(d) for an illustration. Since  $d_{AQ_n}(\mathbf{r}_{k-2}, \mathbf{y}) = 2$ , by Lemma 4, there exists a vertex  $\mathbf{p} \in Nbd_{AQ_n}(\mathbf{r}_{k-2})$  in  $V(AQ_{n-1}^1)$  such that  $d_{AQ_n}(\mathbf{p}, \mathbf{y}) = 1$ . Suppose that  $l_2 = 4$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 3$ . Thus,  $\langle \mathbf{x}, \mathbf{r}_1, \mathbf{p}, \mathbf{y} \rangle$  is a shortest path joining  $\mathbf{x}$  and  $\mathbf{y}$ . By Lemma 1, there exists a vertex  $\mathbf{q} \in Nbd_{AQ_{n-1}^1}(\mathbf{p}) \cap Nbd_{AQ_{n-1}^1}(\mathbf{y}) - \{\mathbf{v}\}$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - \{\mathbf{x}, \mathbf{r}_1, \mathbf{p}, \mathbf{q}, \mathbf{y}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . We set  $P_2$  as  $\langle \mathbf{x}, \mathbf{r}_1, \mathbf{p}, \mathbf{q}, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths. Suppose that  $l_2 = 4$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) = 4$ . Thus,  $P_2 = \langle \mathbf{x}, \mathbf{r}_1, \mathbf{r}_2, \mathbf{p}, \mathbf{y} \rangle$  is a shortest path joining  $\mathbf{x}$  and  $\mathbf{y}$ . By Lemma 3, there exists a hamiltonian path  $P_1$  of  $AQ_n - \{\mathbf{x}, \mathbf{r}_1, \mathbf{r}_2, \mathbf{p}, \mathbf{y}\}$  joining  $\mathbf{u}$  to  $\mathbf{v}$ . Obviously,  $P_1$  and  $P_2$  are the required paths. Suppose that  $5 \leq l_2 \leq 2^{n-2}$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \geq 3$ . Obviously, there exists a vertex  $\mathbf{s}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{u}) - \{\mathbf{x}, \mathbf{r}_{k-2}, \mathbf{y}^h, \mathbf{v}^h\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{s}$  with  $l(R_1) = 2^{n-1} - l_2$ , (2)  $R_2$  is a path joining  $\mathbf{r}_{k-2}$  to  $\mathbf{x}$  with  $l(R_2) = l_2 - 2$ , and (3)

$R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . By Lemma 3, there exists a hamiltonian path  $S$  of  $AQ_{n-1}^1 - \{\mathbf{p}, \mathbf{y}\}$  joining  $\mathbf{s}^h$  to  $\mathbf{v}$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^h, S, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, R_2, \mathbf{r}_{k-2}, \mathbf{p}, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths. Suppose that  $2^{n-2} + 1 \leq l_2 < 2^{n-1} - 1$  except  $2^{n-1} - 3$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \geq 3$ . Obviously, there exists a vertex  $\mathbf{s}$  in  $Nbd_{AQ_{n-1}^0}(\mathbf{u}) - \{\mathbf{x}, \mathbf{r}_{k-2}, \mathbf{y}^h, \mathbf{v}^h\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{s}$  with  $l(R_1) = 2^{n-2} + 1$ , (2)  $R_2$  is a path joining  $\mathbf{r}_{k-2}$  to  $\mathbf{x}$  with  $l(R_2) = 2^{n-2} - 3$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . Again by induction, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{s}^h$  to  $\mathbf{v}$  with  $l(S_1) = 2^{n-1} - l_2 + 2^{n-2} - 4$ , (2)  $S_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{y}$  with  $l(S_2) = l_2 - 2^{n-2} + 2$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^h, S_1, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, R_2, \mathbf{r}_{k-2}, \mathbf{p}, S_2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

**Subcase 5.6:**  $l_2 = 2^{n-1} - 3$  or  $l_2 = 2^{n-1} - 1$  with  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \geq 2$ . Let  $t = 0$  if  $l_2 = 2^{n-1} - 3$  and  $t = 1$  if  $l_2 = 2^{n-1} - 1$ . Obviously, there exist two vertices  $\mathbf{s}$  and  $\mathbf{p}$  in  $AQ_{n-1}^0 - \{\mathbf{u}, \mathbf{x}, \mathbf{v}^n, \mathbf{y}^n\}$ . By induction, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{s}$  with  $l(R_1) = 2^{n-2} - t$ , (2)  $R_2$  is a path joining  $\mathbf{p}$  to  $\mathbf{x}$  with  $l(R_2) = 2^{n-2} + t - 2$ , and (3)  $R_1 \cup R_2$  spans  $AQ_{n-1}^0$ . Similarly, there exist two disjoint paths  $S_1$  and  $S_2$  such that (1)  $S_1$  is a path joining  $\mathbf{s}^n$  to  $\mathbf{v}$  with  $l(S_1) = 2^{n-2} - t$ , (2)  $S_2$  is a path joining  $\mathbf{p}^n$  to  $\mathbf{y}$  with  $l(S_2) = 2^{n-2} + t - 2$ , and (3)  $S_1 \cup S_2$  spans  $AQ_{n-1}^1$ . We set  $P_1$  as  $\langle \mathbf{u}, R_1, \mathbf{s}, \mathbf{s}^n, S_1, \mathbf{v} \rangle$  and  $P_2$  as  $\langle \mathbf{x}, R_2, \mathbf{p}, \mathbf{p}^n, S_2, \mathbf{y} \rangle$ . Obviously,  $P_1$  and  $P_2$  are the required paths.

Thus, Theorem 7 is proved.  $\square$

## 4 Concluding remarks

Now, we make some remarks to illustrate that some in-

teresting properties of augmented cubes are consequences of Theorem 7.

**Remark 1.** The hamiltonian connected property of augmented cubes, proved in [8], states that there exists a hamiltonian path of  $AQ_n$  joining any two different vertices  $\mathbf{u}$  and  $\mathbf{y}$ . Now, we prove that  $AQ_n$  is hamiltonian connected by Theorem 7. Obviously,  $AQ_n$  is hamiltonian connected for  $n = 1$ . Since  $n \geq 2$ , we can choose a pair of adjacent vertices  $\mathbf{v}$  and  $\mathbf{x}$  such that  $\{\mathbf{v}, \mathbf{x}\} \cap \{\mathbf{u}, \mathbf{y}\} = \emptyset$ . By Theorem 7, there are two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{u}$  to  $\mathbf{v}$ , (2)  $P_2$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$ , and (3)  $P_1 \cup P_2$  spans  $AQ_n$ . Obviously,  $\langle \mathbf{u}, P_1, \mathbf{v}, \mathbf{x}, P_2, \mathbf{y} \rangle$  forms a hamiltonian path joining  $\mathbf{u}$  to  $\mathbf{y}$ . Thus,  $AQ_n$  is hamiltonian connected.

**Remark 2.** The panconnected property of  $AQ_n$ , proved in [13], stated that between any two different vertices  $\mathbf{x}$  and  $\mathbf{y}$  of  $AQ_n$  there exists a path  $P_l(\mathbf{x}, \mathbf{y})$  of length  $l$  for any  $d_{AQ_n}(\mathbf{x}, \mathbf{y}) \leq l \leq 2^n - 1$ . Now, we prove that  $AQ_n$  is panconnected by Theorem 7. Obviously,  $AQ_n$  is panconnected for  $n = 1, 2$ . Now, we consider that  $n \geq 3$ .

Suppose that  $l = 2^n - 1$ . By Remark 1,  $AQ_n$  is hamiltonian connected. Obviously, the hamiltonian path of  $AQ_n$  joining  $\mathbf{x}$  and  $\mathbf{y}$  is of length  $2^n - 1$ . Suppose that  $l = 2^n - 2$ . Let  $\mathbf{u}$  be a vertex in  $Nbd_{AQ_n}(\mathbf{y}) - \{\mathbf{x}\}$ . By Lemma 1, there exists a vertex  $\mathbf{v}$  in  $(Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{y})) - \{\mathbf{x}\}$ . By Theorem 7, there exist two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{x}$  to  $\mathbf{u}$  with  $l(P_1) = 2^n - 3$ , (2)  $P_2$  is a path joining  $\mathbf{y}$  to  $\mathbf{v}$  with  $l(P_2) = 1$ , and (3)  $P_1 \cup P_2$  spans  $AQ_n$ . Obviously,  $\langle \mathbf{x}, P_1, \mathbf{u}, \mathbf{y} \rangle$  is a path of length  $2^n - 2$  joining  $\mathbf{x}$  to  $\mathbf{y}$ . Suppose that  $l = 2^n - 3$ . We can find two adjacent vertices  $\mathbf{u}$  and  $\mathbf{v}$  such that  $\{\mathbf{u}, \mathbf{v}\} \cap \{\mathbf{x}, \mathbf{y}\} = \emptyset$ . By Theorem 7, there exist two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$  with  $l(P_1) = 2^n - 3$ , (2)  $P_2$  is a path joining  $\mathbf{u}$  to  $\mathbf{v}$  with  $l(P_2) = 1$ , and (3)

$P_1 \cup P_2$  spans  $AQ_n$ . Obviously,  $P_1$  is a path of length  $2^n - 3$  joining  $\mathbf{x}$  to  $\mathbf{y}$ . Suppose that  $l \leq 2^n - 4$ . By Lemma 2, there exist two vertices  $\mathbf{u}$  and  $\mathbf{v}$  such that  $d_{AQ_n}(\mathbf{u}, \mathbf{v}) = 2$ ,  $\{\mathbf{x}, \mathbf{y}\} \neq Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{v})$ , and  $\{\mathbf{u}, \mathbf{v}\} \neq Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . By Theorem 7, there exist two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$  with  $l(P_1) = l$ , (2)  $P_2$  is a path joining  $\mathbf{u}$  to  $\mathbf{v}$  with  $l(P_2) = 2^n - 2 - l$ , and (3)  $P_1 \cup P_2$  spans  $AQ_n$ . Obviously,  $P_1$  is a path of length  $l$  joining  $\mathbf{x}$  to  $\mathbf{y}$ . Thus,  $AQ_n$  is panconnected.

**Remark 3.** The edge-pancyclic property property of  $AQ_n$  stated that for any edge  $e = (\mathbf{x}, \mathbf{y})$  and for any  $3 \leq l \leq 2^n$ , there exists a cycle of length  $l$  containing  $e$  if  $n \geq 2$ . We prove that  $AQ_n$  is edge-pancyclic by Theorem 7. Obviously,  $AQ_n$  is edge-pancyclic for  $n = 2$ . Thus, we consider that  $n \geq 3$ .

Suppose that  $l = 3$ . By Lemma 1, there exists  $\mathbf{u} \in Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ . Obviously,  $\langle \mathbf{x}, \mathbf{y}, \mathbf{u}, \mathbf{x} \rangle$  forms a cycle of length three containing  $e$ . Now, we consider that  $l = 2^n$  and  $l = 2^n - 1$ . By Lemma 1, there exists  $\mathbf{v} \in (Nbd_{AQ_n}(\mathbf{u}) \cap Nbd_{AQ_n}(\mathbf{y})) - \{\mathbf{x}\}$ . By Theorem 7, there exist two disjoint paths  $P_1$  and  $P_2$  such that (1)  $P_1$  is a path joining  $\mathbf{x}$  to  $\mathbf{u}$  with  $l(P_1) = 2^n - 3$ , (2)  $P_2$  is a path joining  $\mathbf{v}$  to  $\mathbf{y}$  with  $l(P_2) = 1$ , and (3)  $P_1 \cup P_2$  spans  $AQ_n$ . Obviously,  $\langle \mathbf{x}, P_1, \mathbf{u}, \mathbf{v}, \mathbf{y}, \mathbf{x} \rangle$  forms a cycle of length  $2^n$  containing  $e$  and  $\langle \mathbf{x}, P_1, \mathbf{u}, \mathbf{y}, \mathbf{x} \rangle$  forms a cycle of length  $2^n - 1$  containing  $e$ . Suppose  $l = 2^n - 2$ . By Theorem 7, there exist two disjoint paths  $Q_1$  and  $Q_2$  such that (1)  $Q_1$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$  with  $l(Q_1) = 2^n - 3$ , (2)  $Q_2$  is a path joining  $\mathbf{u}$  to  $\mathbf{v}$  with  $l(Q_2) = 1$ , and (3)  $Q_1 \cup Q_2$  spans  $AQ_n$ . Obviously,  $\langle \mathbf{x}, Q_1, \mathbf{y}, \mathbf{x} \rangle$  forms a cycle of length  $2^n - 2$  containing  $e$ . Suppose that  $4 \leq l \leq 2^n - 3$ . By Lemma 2, there exists two vertices  $\mathbf{p}$  and  $\mathbf{q}$  of  $AQ_n$  such that  $d_{AQ_n}(\mathbf{p}, \mathbf{q}) = 2$ ,  $\{\mathbf{x}, \mathbf{y}\} \neq Nbd_{AQ_n}(\mathbf{p}) \cap Nbd_{AQ_n}(\mathbf{q})$ , and  $\{\mathbf{p}, \mathbf{q}\} \neq Nbd_{AQ_n}(\mathbf{x}) \cap Nbd_{AQ_n}(\mathbf{y})$ .

By Theorem 7, there exist two disjoint paths  $R_1$  and  $R_2$  such that (1)  $R_1$  is a path joining  $\mathbf{x}$  to  $\mathbf{y}$  with  $l(R_1) = l - 1$ , (2)  $R_2$  is a path joining  $\mathbf{u}$  to  $\mathbf{v}$  with  $l(R_2) = 2^n - l - 1$ , and (3)  $R_1 \cup R_2$  spans  $AQ_n$ . Obviously,  $\langle \mathbf{x}, R_1, \mathbf{y}, \mathbf{x} \rangle$  forms a cycle of length  $l$  containing  $e$ .

## References

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, North-Holland, New York, 1980.
- [2] S.A. Choudum, V. Sunitha, Augmented cubes, Networks 40 (2) (2002) 71–84.
- [3] J.-M. Chang, J.-S. Yang, Y.-L. Wang, Y. Cheng, Panconnectivity, fault-tolerant hamiltonicity and hamiltonian connectivity in alternating group graphs, Networks 44 (2004) 302–310.
- [4] J. Fan, X. Jia, X. Lin, Complete path embeddings in crossed cubes, Information Science, 176 (2006) 3332–3346.
- [5] J. Fan, X. Jia, X. Lin, Optimal embeddings of paths with various lengths in twisted cubes, IEEE Trans. Parallel and Distributed Systems, 18, (2007) 511–521.
- [6] A. Germa, M.C. Heydemann, D. Sotteau, Cycles in cube-connected cycles graph, Discrete Appl. Math. 83 (1998) 135–155.
- [7] T.Y. Ho, C.K. Lin, Jimmy J.M. Tan, L.H. Hsu, The super spanning connectivity of augmented cubes, accepted by *Ars Combinatorica*.
- [8] H.C. Hsu, L.C. Chiang, J.J.M. Tan, L.H. Hsu, Fault hamiltonicity of augmented cubes, Parallel Comput. 31 (2005) 130–145.

- [9] H.C. Hsu, P.L. Lai, C.H. Tsai, Geodesic pancyclicity and balanced pancyclicity of augmented cubes, *Information Processing Letters* 101 (2007) 227–232.
- [10] S.C. Hwang, G.H. Chen, Cycles in butterfly graphs, *Networks* 35 (2) (2000) 161–171.
- [11] F.T. Leighton, *Introduction to Parallel Algorithms and Architectures: Arrays · Trees · Hypercubes*, Morgan Kaufmann Publishers, San Mateo, CA, 1992.
- [12] T.K. Li, C.H. Tsai, Jimmy J.M. Tan, L.H. Hsu, Bipanconnected and edge-fault-tolerant bipancyclic of hypercubes, *Information Processing Letters*, 87 (2003) 107–110.
- [13] M. Ma, G. Liu, J.M. Xu, Panconnectivity and edge-fault-tolerant pancyclicity of augmented cubes, *Parallel Computing* 33 (2007) 35–42.
- [14] M. Ma, J.M. Xu, Panconnectivity of locally twisted cubes, *Applied Mathematics Letters*, 19, (2006) 673–677.