

# On the Two-Equal-Disjoint Path Cover Problem of Twisted Cubes

Pao-Lien Lai

Hong-Chun Hsu

Department of Computer Science  
and Information Engineering  
National Dong Hwa University,  
Shoufeng, Hualien, Taiwan 97401, R.O.C.  
baolein@mail.ndhu.edu.tw

Department of Medical Informatics  
Tzu Chi University,  
Hualien, Taiwan 970, R.O.C.  
hchsu@mail.tcu.edu.tw

## Abstract

Embedding of paths have attracted much attention in the parallel processing. Many-to-many communication is one of the most central issues in various interconnection networks. A graph  $G$  is globally two-equal-disjoint path coverable if for any two distinct pairs of vertices  $(u, v)$  and  $(x, y)$  of  $G$ , there exist two disjoint paths  $P_{uv}$  and  $P_{xy}$  satisfied that (1)  $P_{uv}$  joins  $u$  to  $v$  and  $P_{xy}$  joins  $x$  to  $y$ , (2)  $|P_{uv}| = |P_{xy}|$ , and (3)  $V(P_{xy} \cup P_{xy}) = V(G)$ . In this paper, we prove that  $TQ_n$  is globally 2-equal-disjoint path coverable for  $n \geq 5$ .

*Keywords:* Interconnection network; Twisted cube; disjoint path;  $k$ -equal-disjoint path cover, 2-equal-disjoint path coverable.

## 1 Introduction

For the graph definition and notation we follow [2].  $G = (V, E)$  is a graph if  $V$  is a finite set and  $E$  is a subset of  $\{(a, b) \mid (a, b) \text{ is an unordered pair of } V\}$ . We say that  $V$  is the *vertex set* and  $E$  is the *edge set*. A *path* of length  $k$  from  $x$  to  $y$  is a finite sequence of distinct vertices  $\langle v_0, v_1, v_2, \dots, v_k \rangle$ , where  $x = v_0$ ,  $y = v_k$ , and  $(v_{i-1}, v_i) \in E$  for all  $1 \leq i \leq k$ . For convenience, we use the sequence  $\langle v_0, \dots, v_i, P, v_j, \dots, v_k \rangle$ , where  $P = \langle v_i, v_{i+1}, \dots, v_j \rangle$  to denote the path  $\langle v_0, v_1, v_2, \dots, v_k \rangle$ . Note that it is possible that the path  $P$  has length 0. We can also write the path  $\langle v_0, v_1, v_2, \dots, v_k \rangle$  as  $\langle v_0, P_1, v_i, v_{i+1}, \dots, v_j, P_2, v_t, \dots, v_k \rangle$ , where  $P_1$  is the path  $\langle v_0, v_1, \dots, v_i \rangle$  and  $P_2$  is the path  $\langle v_j, v_{j+1}, \dots, v_t \rangle$ .

---

\*Correspondence to: Assistant Professor Pao-Lien Lai, Department of Computer Science and Information Engineering, National Dong Hwa University, Shoufeng, Hualien, Taiwan 97401, R.O.C. e-mail:baolein@mail.ndhu.edu.tw

This work was supported in part by the National Science Council of the Republic of China under Contract NSC 95-2221-E-259-045.

We use  $d(u, v)$  to denote the distance between  $u$  and  $v$ , i.e., the length of the shortest path joining  $u$  and  $v$ .

A path is a *Hamiltonian path* if it contains all vertices of  $G$ . A graph  $G$  is *Hamiltonian connected* if there exists a Hamiltonian path joining any two distinct vertices. A *cycle* is a path (except the first vertex is the same as the last vertex) containing at least three vertices. A cycle of  $G$  is a *Hamiltonian cycle* if it contains all vertices. A graph is *Hamiltonian* if it has a Hamiltonian cycle.

Finding node-disjoint paths is one of the important issues of routing among nodes in various interconnection networks. Node-disjoint (abbreviated as disjoint) paths can be used to avoid communication congestion and provide parallel paths for an efficient data routing among nodes. Moreover, multiple disjoint paths can be more fault-tolerant of node or link failures and greatly enhance the transmission reliability. Disjoint paths generally fall into three categories: one-to-one, one-to-many, and many-to-many. The one-to-one disjoint path is built with one source and one destination. The one-to-many disjoint paths like a tree structure, they contain one source and many distinct destination nodes. The many-to-many disjoint paths involve  $k, k \geq 1$ , disjoint paths with  $k$  pairs distinct source and destination nodes.

A *disjoint path cover* in a graph  $G$  is to find disjoint paths containing all the vertices in  $G$ . For an embedding of linear arrays in a network, the cover implies every node can be participated in a pipeline computation. One-to-one disjoint path covers in recursive circulants [8] and one-to-many disjoint path covers in some hypercube-like interconnection networks [9] were studied. The many-to-many  $k$ -disjoint path cover is proposed by Park etc. in [10]. In this paper, we call such many-to-many  $k$ -disjoint path cover (abbreviated as  $k$ -disjoint path cover) as many-to-many  $k$ -equal-disjoint path cover (abbreviated as  $k$ -equal-disjoint path cover) that  $k$  disjoint paths have same lengths. The  $k$  disjoint paths with equal length implies that the parallel processing of  $k$  pipeline is guaranteed accurately. Furthermore, a graph is called globally  $k$ -equal-

disjoint path coverable if there exists a  $k$ -equal-disjoint path cover for any  $k$  distinct source-destination pairs.

An  $n$ -dimensional Twisted cube,  $TQ_n$  [3], is a variation of hypercube. It has  $2^n$  vertices and each vertex has the same degree  $n$ . The difference is that it diverts some edges which are called *twisted edge* and these edge reduce the distance between vertices.

Though some topological properties of Twisted cubes have been studied in the literature [1, 4, 5, 7, 11, 12, 13]. In this paper, we prove that the Twisted cube is globally two-equal-disjoint path coverable. In next section, we give the definition of two-equal-disjoint path coverable problem and Twisted Cubes. Then we prove that the Twisted cube is globally two-equal-disjoint path coverable in the section 3. In the final section, we give the conclusion.

## 2 Preliminary

In this section, we will first give the definition of globally two-equal-disjoint path coverable problem of a graph  $G$ , and then we will give the relevant definitions in graph theory and the definition of the Twisted cubes.

**Definition 1** A graph  $G$  is  $(u, v, x, y)$ -two-equal-disjoint path coverable if there are two disjoint paths  $P_{uv}$  and  $P_{xy}$  such that  $P_{uv}$  joins the vertices  $u$  to  $v$ ,  $P_{xy}$  joins the vertices  $x$  to  $y$ , and  $V(P_{uv} \cup P_{xy}) = V(G)$ .

**Definition 2** A graph  $G$  is globally two-equal-disjoint path coverable if for any two distinct pairs of vertices  $(u, v)$  and  $(x, y)$ , the  $(u, v, x, y)$ -two-equal-disjoint path cover exists.

To define the Twisted cubes, as proposed by Hilbers [3], the term so called "parity" is introduced.

**Definition 3** The *parity*  $P_j(u) = u_j \oplus u_{j-1} \oplus \dots \oplus u_1 \oplus u_0$  is a function using exclusive-or,  $\oplus$ .

A 2-bit in  $u$  is a pair of adjacent bits where the larger index is even, for example  $u_2u_1$ . And the  $i$ -th 2-bit is  $u_{2i}u_{2i-1}$  for  $i \geq 1$ . But the 0-th 2-bit is just  $u_0$ . The following is the recursive definition of the  $n$ -dimensional Twisted cube  $TQ_n$ .

**Definition 4** [3] An  $n$ -dimensional twisted cube for  $n = 2k + 1$ , denoted  $TQ_n$ , is a graph  $G(V, E)$ , where  $V = \{v | v \in \{0, 1\}^n\}$  and  $E = \{(x, y) | x, y \in V, \text{ satisfying}$

- (1)  $x_{2i}x_{2i-1} = \bar{y}_{2i}y_{2i-1}$  or  $y_{2i}\bar{y}_{2i-1}$ ; with  $P_{2i-2}(x) = 1$ , or
- $x_{2i}x_{2i-1} = \bar{y}_{2i}y_{2i-1}$  or  $\bar{y}_{2i}\bar{y}_{2i-1}$ ; with  $P_{2i-2}(x) = 0$  for some 2-bit  $i$ ; and

- (2)  $x_{2j}x_{2j-1} = y_{2j}y_{2i-1}$  for any other 2-bits  $j \neq i$ .

A *matching*  $M$  of a graph  $G$  is a set of pairwise disjoint edges.  $M$  is a perfect matching if each vertex of  $G$  belongs to some edge in  $M$ . From the definition, we have the following lemma.

**Lemma 1** [6]

Both the subgraph induced by  $TQ_{n-2}^{0,0} \cup TQ_{n-2}^{1,0}$  and the subgraph induced by  $TQ_{n-2}^{0,1} \cup TQ_{n-2}^{1,1}$  are isomorphic to  $TQ_{n-2} \times K_2$  where  $K_2$  is the complete graph with two vertices. Moreover, the edges joining  $TQ_{n-2}^{0,0} \cup TQ_{n-2}^{1,0}$  and  $TQ_{n-2}^{0,1} \cup TQ_{n-2}^{1,1}$  form a perfect matching of  $TQ_n$ .

Let  $G$  and  $H$  be two graphs having the same number of vertices.  $G \oplus_M H$  denotes a graph which has copies of  $G$  and  $H$  connected by a matching  $M$ . Let  $TQ_{n+1}^0$  and  $TQ_{n+1}^1$  be the subgraphs induced by  $V(TQ_n^{0,0}) \cup V(TQ_n^{1,0})$  and  $V(TQ_n^{0,1}) \cup V(TQ_n^{1,1})$ , respectively. Then by Lemma 1, both of  $TQ_{n+1}^0$  and  $TQ_{n+1}^1$  are isomorphic to  $TQ_n \times K_2$ , and  $TQ_{n+1}^0 \oplus_M TQ_{n+1}^1$  is isomorphic to  $TQ_{n+2}$  for a specific matching  $M$ . In addition,  $TQ_n \times K_2$  has two copies of  $TQ_n$ , and we use  $TQ_n^0$  and  $TQ_n^1$  to denote them, respectively. For convenience of discussion, we add 0 to every vertex  $v \in V(TQ_n^0)$  and 1 to every vertex  $u \in V(TQ_n^1)$ , respectively, as the leading bits. As a result, each vertex of  $TQ_n \times K_2$  is represented by a binary string of length  $n + 1$ . These notations are used extensively throughout this paper.

We then introduce two important fault Hamiltonian results for proving the main theorem in the next section of this paper. A graph  $G$  is  $k$ -fault Hamiltonian connected if for any faulty set  $F \subset V(G) \cup E(G)$  such that  $|F| \leq k$ ,  $G - F$  is still Hamiltonian connected.

**Lemma 2** [6]  $TQ_n$  is  $n - 3$  fault Hamiltonian connected.

**Lemma 3** [6] For  $n \geq 3$  and  $i \in \{0, 1\}$ ,  $TQ_n^{0,i} \cup TQ_n^{1,i}$  is  $(n - 1)$ -Hamiltonian and  $(n - 2)$ -Hamiltonian connected

## 3 Twisted cube is globally two-disjoint equal path coverable

As a starting point we present lemmas 4 and 5 which establish the base case of Theorem 1.

**Lemma 4**  $TQ_3$  is not globally two-equal-disjoint path coverable.

**Proof.** To prove this lemma, we give a counterexample. Given two pair of vertices 0, 1 and 2, 4,  $TQ_3$  is not (0, 1, 2, 4)-two-equal-disjoint path coverable.  $\square$

**Lemma 5**  $TQ_5$  is globally two-equal-disjoint path coverable.

**Proof.** To prove this case is very tedious. With long and detail discussion, we have completed theoretical proof for  $TQ_5$ . Nevertheless, we do not present it in this paper for reducing complexity. However, we can also verify this small case directly using computer.  $\square$

Next we show  $TQ_n \times K_2$  has the same result if  $TQ_n$  is globally two-equal-disjoint path coverable.

**Lemma 6** Let  $n \geq 5$ .  $TQ_n \times K_2$  is also globally two-equal-disjoint path coverable if  $TQ_n$  is globally two-equal-disjoint path coverable.

**Proof.** Let  $TQ_{n+1} = TQ_n \times K_2$  and let  $TQ_n^0$  and  $TQ_n^1$  be the two components of  $TQ_{n+1}$ . Let  $(u, v)$  and  $(x, y)$  be any two distinct vertex pairs of  $TQ_{n+1}$ . Herein, we want to establish two disjoint paths  $P_{uv}$ , and  $P_{xy}$  with equal length  $2^n - 1$ . The proof consists of four cases as follows:

**Case 1:** All four end vertices belong to the same  $TQ_n^i$ ,  $i = 0, 1$ .

Without loss of generality, let  $u, v, x, y \in V(TQ_n^0)$ . By hypothesis, there exist two equal disjoint paths  $P_{uv}^0$  and  $P_{xy}^0$  of length  $2^{n-1} - 1$  in  $TQ_n^0$ . Let  $t$  (resp.  $w$ ) be the neighbor of  $u$  (resp.  $x$ ) on path  $P_{uv}^0$  (resp.  $P_{xy}^0$ ). Let  $P_{uv}^0 = (u, t, P_{tv}^0)$  and  $P_{xy}^0 = (x, w, P_{wy}^0)$ . Let  $u^1, t^1, x^1$ , and  $w^1$  be the neighbors of  $u, t, x$ , and  $w$  in  $TQ_n^1$ , respectively. Then, there are also two equal disjoint paths  $P_{u^1t^1}^1$  and  $P_{x^1w^1}^1$  of length  $2^{n-1} - 1$  in  $TQ_n^1$ . Let  $P_{uv} = (u, u^1, P_{u^1t^1}^1, t^1, t, P_{tv}^0, v)$  and  $P_{xy} = (x, x^1, P_{x^1w^1}^1, w^1, w, P_{wy}^0, y)$ . Hence,  $P_{uv}$  and  $P_{xy}$  are two equal disjoint paths of length  $2^n - 1$  in  $TQ_n \times K_2$ .

**Case 2:** Three end vertices belong to  $TQ_n^i$  and another end vertex belongs to  $TQ_n^{1-i}$ ,  $i = 0, 1$ .

Without loss of generality, let  $u, v, x \in V(TQ_n^0)$  and  $y \in V(TQ_n^1)$ . Let  $y^0$  be the neighbors of  $y$  in  $TQ_n^0$ . Let  $w$  be a vertex in  $TQ_n^0$  and  $w \notin \{u, v, x, y^0\}$ . By hypothesis, there exist two equal disjoint paths  $P_{uv}^0$  and  $P_{xw}^0$  of length  $2^{n-1} - 1$  in  $TQ_n^0$ . Let  $(s, t) \in P_{uv}^0$  and  $y^0 \notin \{s, t\}$ . Let  $P_{uv}^0 = \langle u, P_{us}^0, s, t, P_{tv}^0, v \rangle$ .

Let  $s^1, t^1, w^1$  be the neighbors of  $s, t, w$  in  $TQ_n^1$ , respectively. Note that there exist two equal disjoint paths  $P_{s^1t^1}^1$  and  $P_{w^1y^0}^0$  of length  $2^{n-1} - 1$  in  $TQ_n^1$ . Let  $P_{uv} = \langle u, P_{us}^0, s, s^1, P_{s^1t^1}^1, t^1, t, P_{tv}^0, v \rangle$  and  $P_{xy} = \langle x, P_{xw}^0, w, w^1, P_{w^1y^0}^0, y \rangle$ . Then,  $P_{uv}$  and  $P_{xy}$  are two equal disjoint paths of length  $2^n - 1$  in  $TQ_n \times K_2$ .

**Case 3:**  $u, v$  belong to  $TQ_n^i$  and  $x, y$  belong to  $TQ_n^{1-i}$ ,  $i = 0, 1$ .

Without loss of generality, let  $u, v \in V(TQ_n^0)$  and  $x, y \in V(TQ_n^1)$ . By Lemma 2, there exist two paths  $P_{uv}$  and  $P_{xy}$  of equal length  $2^n - 1$  in  $TQ_n^0$  and  $TQ_n^1$ , respectively.

**Case 4:**  $u, x$  (or  $u, y$ ) belong to  $TQ_n^i$  and  $v, y$  (or  $v, x$ ) belong to  $TQ_n^{1-i}$ ,  $i = 0, 1$ .

Without loss of generality, let  $u, x \in V(TQ_n^0)$  and  $v, y \in V(TQ_n^1)$ . Let  $v^0, y^0$  be the neighbors of  $v, y$  in  $TQ_n^0$ , respectively. Let  $s, w$  be two vertices except  $u, x, v^0, y^0$  in  $TQ_n^0$  and let  $s^1, w^1$  be the neighbors of  $s, w$  in  $TQ_n^1$ , respectively. Let  $P_{us}$  and  $P_{xw}$  be two disjoint paths of length  $2^{n-1} - 1$  in  $TQ_n^0$ . Let  $P_{s^1v}$  and  $P_{w^1y}$  be two disjoint paths of length  $2^{n-1} - 1$  in  $TQ_n^1$ . Let  $P_{uv} = \langle u, P_{us}, s, s^1, P_{s^1v}, v \rangle$  and  $P_{xy} = \langle x, P_{xw}, w, w^1, P_{w^1y}, y \rangle$ . Then  $P_{uv}$  and  $P_{xy}$  be two disjoint paths of lengths  $2^n - 1$  in  $TQ_n \times K_2$ .  $\square$

Next we formally show the main result that  $TQ_n$ ,  $n \geq 5$ , is globally two-disjoint equal path coverable.

**Theorem 1** Twisted cube,  $TQ_n$ , is globally two-equal-disjoint path coverable for  $n \geq 5$ .

**Proof.** We prove this theorem by induction on  $n$ . The base case is  $TQ_5$ . With Lemma 5, the base case holds. By induction hypothesis, we can assume that  $TQ_n$  is globally two-equal-disjoint path coverable. Now, we need to show that  $TQ_{n+2}$  is also globally two-equal-disjoint path coverable. Let  $u, v$  and  $x, y$  be two distinct source-destination pairs of  $TQ_{n+2}$ . In the following, we establish two disjoint paths  $P_{uv}, P_{xy}$  of length  $2^{n+1} - 1$ . Herein, we divide the proof into two cases according to which subgraphs,  $TQ_n^{0,i} \cup TQ_n^{1,i}$ , the four vertices exactly belong to as follows.

**Case 1:** The four vertices  $u, v, x, y$  exactly belong to one of the two subgraphs  $TQ_n^{0,i} \cup TQ_n^{1,i}$  and  $TQ_n^{1,1-i} \cup TQ_n^{0,1-i}$ .

Without loss of generality, let  $i = 0$ . By Lemma 6, there exist two disjoint paths, denoted as  $P_{uv}^0$  and  $P_{xy}^0$ , of length  $2^n - 1$  in  $TQ_n^{0,0} \cup TQ_n^{1,0}$ . Let  $s$  and  $w$  be the neighbor of  $u$  and  $x$  on paths  $P_{uv}^0$  and  $P_{xy}^0$ , respectively. Let  $P_{uv}^0 = \langle u, s, P_{sv}^0, v \rangle$  and  $P_{xy}^0 = \langle x, w, P_{wy}^0, y \rangle$ . Let  $u^1, s^1, x^1, w^1$  be the neighbors of vertices  $u, s, x, w$  in  $TQ_n^{0,1} \cup TQ_n^{1,1}$ , respectively. By Lemma 6, there also exist two disjoint paths, denoted as  $P_{u^1s^1}^1$  and  $P_{x^1w^1}^1$ , of length  $2^n - 1$  in  $TQ_n^{0,1} \cup TQ_n^{1,1}$ . Hence,  $P_{uv} = \langle u, u^1, P_{u^1s^1}^1, s^1, s, P_{sv}^0, v \rangle$  and  $P_{xy} = \langle x, x^1, P_{x^1w^1}^1, w^1, w, P_{wy}^0, y \rangle$  are two disjoint paths of length  $2^{n+1} - 1$  in  $TQ_{n+2}$ .

**Case 2:** The four vertices  $u, v, x, y$  exactly belong to the two subgraphs  $TQ_n^{0,0} \cup TQ_n^{1,0}$  and  $TQ_n^{0,1} \cup TQ_n^{1,1}$ .

Herein, we in advanced divide this case into three sub-cases as follows.

**Subcase 2.1:** Three end vertices belong to  $TQ_n^{0,i} \cup TQ_n^{1,i}$  and another vertex belongs to  $TQ_n^{0,1-i} \cup TQ_n^{1,1-i}$ .

Without loss of generality, let  $u, v, x \in V(TQ_n^{0,0}) \cup V(TQ_n^{1,0})$  and  $y \in V(TQ_n^{0,1}) \cup V(TQ_n^{1,1})$ . Let vertex  $y^0 \in V(TQ_n^{0,0}) \cup V(TQ_n^{1,0})$  be a neighbor of  $y$ . Let  $w$  be a neighbor of  $x$  in  $TQ_n^{0,0} \cup TQ_n^{1,0}$  except  $u, v$  and  $y^0$ . By Lemma 3, there exist path  $P_{uv}^0$  of length  $2^{n+1} - 3$  in  $TQ_n^{0,0} \cup TQ_n^{1,0}$  except  $w$  and  $x$ . Let  $(s, t)$  be an edge on path  $P_{uv}^0$  with  $y^0 \notin \{s, t\}$ . Let  $P_{uv}^0 = \langle u, P_{us}^0, s, t, P_{tv}^0, v \rangle$ . Let vertices  $s^1, t^1, w^1 \in V(TQ_n^{0,1}) \cup V(TQ_n^{1,1})$  be the neighbors of  $s, t, w$ , respectively. By Lemma 3, there exist path  $P_{w^1y}$  of length  $2^{n+1} - 3$  in  $TQ_n^{0,1} \cup TQ_n^{1,1}$  except  $s^1$  and  $t^1$ .

Let  $P_{uv} = \langle u, P_{us}^0, s, s^1, t^1, t, P_{tv}^0, v \rangle$  and  $P_{xy} = \langle x, w, w^1, P_{w^1y}, y \rangle$ . Clearly,  $P_{uv}$  and  $P_{xy}$  are two disjoint paths of length  $2^{n+1} - 1$ .

**Subcase 2.2:** Each pair has one vertex in  $TQ_n^{0,i} \cup TQ_n^{1,i}$  and another vertex in  $TQ_n^{0,1-i} \cup TQ_n^{1,1-i}$ .

Without loss of generality, let  $u, x \in V(TQ_n^{0,0}) \cup V(TQ_n^{1,0})$  and  $v, y \in V(TQ_n^{0,1}) \cup V(TQ_n^{1,1})$ . Let  $u^1, x^1 \in V(TQ_n^{0,1}) \cup V(TQ_n^{1,1})$  and  $v^0, y^0 \in V(TQ_n^{0,0}) \cup V(TQ_n^{1,0})$  be the neighbors of  $u, v$  and  $v, y$ , respectively. Let  $s$  and  $w$  be two vertices in  $TQ_n^{0,0} \cup TQ_n^{1,0}$  except  $u, x, v^0, y^0$ . Let  $s^1$  and  $w^1$  be the neighbors of  $s$  and  $w$ , respectively, in  $TQ_n^{0,1} \cup TQ_n^{1,1}$ .

By Lemma 6, there exist two disjoint paths  $P_{us}^0, P_{xw}^0$  in  $TQ_n^{0,0} \cup TQ_n^{1,0}$  and exist two disjoint paths  $P_{s^1v}^1, P_{w^1y}^1$  in  $TQ_n^{0,1} \cup TQ_n^{1,1}$  of length  $2^n - 1$ , respectively. Then  $P_{uv} = \langle u, P_{us}^0, s, s^1, P_{s^1v}^1, v \rangle$  and  $P_{xy} = \langle x, P_{xw}^0, w, w^1, P_{w^1y}^1, y \rangle$  are two disjoint paths of length  $2^{n+1} - 1$ .

**Subcase 2.3:** One pair belongs to  $TQ_n^{0,i} \cup TQ_n^{1,i}$  and another pair belongs to  $TQ_n^{0,1-i} \cup TQ_n^{1,1-i}$ .

Without loss of generality, let  $u, v \in V(TQ_n^{0,0}) \cup V(TQ_n^{1,0})$  and  $x, y \in V(TQ_n^{0,1}) \cup V(TQ_n^{1,1})$ . By Lemma 3, there exist two disjoint paths  $P_{uv}$  in  $TQ_n^{0,0} \cup TQ_n^{1,0}$  and  $P_{xy}$  in  $TQ_n^{0,1} \cup TQ_n^{1,1}$ , respectively, of length  $2^{n+1} - 1$ .  $\square$

## 4 Conclusion

In this paper, we discussed the two-equal-disjoint path coverable problem and proved that Twisted Cubes  $TQ_n$  are globally two-equal-disjoint path coverable for  $n \geq 5$ . The globally two-equal-disjoint path coverable problem

is an extension of Hamiltonian connected problem. We can see Hamiltonian connected problem as globally one-path coverable problem, and then we extended this property to globally two-equal-disjoint path coverable. This work may help to discuss the many-to-many disjoint path coverable problem.

## References

- [1] S. Abraham and K. S. Padmanabhan, "The twisted cube topology for multiprocessors: a study in network asymmetry," *Journal of Parallel and Distributed Computing*, North Holland, vol. 13, pp. 104-110, 1991.
- [2] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*, North Holland, New York, 1980.
- [3] P.A.J. Hilbers, R.J.M. Koopman, and J.L.A. Van de Snepscheut, "The twisted cube," *Parallel Architectures Languages Europe, Lecture Notes in Computer Science*, (1987), pp. 152-158.
- [4] S. Y. Hsieh, G. H. Chen, and C. W. Ho, "Fault-Free Hamiltonian ring embedding in faulty arrangement graphs," *IEEE Trans. Parallel Distrib. Systems*, vol. 10, pp. 223V237, 1999.
- [5] W. T. Huang, J. M. Tan, C. N. Hung, and L. H. Hsu, "Token ring embedding in faulty twisted cubes," in *Proceeding of the 2nd Inter. Conf. Parallel Sys. PCS'99*, pp. 1V10, 1999.
- [6] W.T. Huang, J.M. Tan, C.N. Hung, and L.H. Hsu, "Fault-Tolerant Hamiltonicity of Twisted Cubes," *Journal of Parallel and Distributed Computing*, Vol.1 62, pp. 591-604, 2002.
- [7] S. Latifi, S. Q. Zheng, and N. Bagherzadeh, "Optimal ring embedding in hypercubes with faulty links," *Proc. IEEE Symp. Fault-Tolerant Comput.*, vol. 42, pp. 178V184, 1992.
- [8] J. H. Park, "One-to-one disjoint path covers in recursive circulants," *Journal of KISS*, 30(12), pp. 691-698, 2003.
- [9] J. H. Park, "One-to-many disjoint path covers in a graph with faulty elements," in *Proc. of the International Computing and Combinatorics Conference COCOON 2004*, pp. 392- 401, 2004.
- [10] J. H. Park, "Many-to-many disjoint path covers in a graph with faulty elements," *International Symposium on Algorithms and Computation, ISAAC 2004*, pp. 742-753, 2004.

- [11] R. A. Rowley and B. Bose, "Fault-tolerant ring embedding in de-Bruijn networks," *IEEE Trans. Comp.*, vol. 12, pp. 1480V1486, 1993.
- [12] T. Y. Sung, C. Y. Lin, Y. C. Chuang, and L. H. Hsu, "Fault tolerant token ring embedding in double loop networks," *Inform. Process. Lett.*, vol. 66, pp. 201V207, 1998.
- [13] Y. C. Tseng, S. H. Chang, and J. P. Sheu, "Fault-tolerant ring embedding in a Star graph with both link and node failures," *IEEE Trans. Parallel. Distrib. Systems*, 1185V1195, 1997.